

Evanston COVID-19 Community Risk Level as of 6/15/2022

| | |
|---|---------------|
| New Cases per 100,000 Population (7-Day Total) | 258.61 |
|---|---------------|

| Indicator | Metric | Evanston Community Risk Level |
|---|--------|-------------------------------|
| New COVID-19 admissions per 100,000 population (7-day total) | 5.12 | MEDIUM |
| Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average) | 1.98% | MEDIUM |

Note: Use prevention strategies from the higher of the two

The Evanston COVID-19 Community Risk Level is based on the following table which was created by the [Centers for Disease Control and Prevention](https://www.cdc.gov):

| COVID-19 Community Levels (Use the Higher Level) | | | | |
|--|---|---------|--------------|---------|
| New COVID-19 cases per 100,000 people in the past 7 days | Indicators | Low | Medium | High |
| Fewer than 200 | New COVID-19 admissions per 100,000 population (7-day total) | < 10.0 | 10.0 - 19.9 | ≥ 20.0 |
| | Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average) | < 10.0% | 10.0 - 14.9% | ≥ 15% |
| 200 or more | New COVID-19 admissions per 100,000 population (7-day total) | N/A | < 10 | ≥ 10 |
| | Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average) | N/A | < 10.0% | ≥ 10.0% |

| COVID-19 Community Level | Individual and household-level prevention behaviors | Community-level prevention strategies |
|--------------------------|--|--|
| Low | <ul style="list-style-type: none"> • Stay up to date with COVID-19 vaccines and boosters • Maintain improved ventilation | <ul style="list-style-type: none"> • Distribute and administer vaccines to achieve high community vaccination coverage and ensure health equity |

| | | |
|--|--|--|
| | <p>throughout indoor spaces when possible</p> <ul style="list-style-type: none"> ● Follow CDC recommendations for isolation and quarantine, including getting tested if you are exposed to COVID-19 or have symptoms of COVID-19 ● If you are immunocompromised or high risk for severe disease <ul style="list-style-type: none"> ○ Have a plan for rapid testing if needed (e.g., having home tests or access to testing) ○ Talk to your healthcare provider about whether you are a candidate for treatments like oral antivirals, PrEP, and monoclonal antibodies | <ul style="list-style-type: none"> ● Maintain improved ventilation in public indoor spaces ● Ensure access to testing, including through point-of-care and at-home tests for all people <ul style="list-style-type: none"> ○ Communicate with organizations and places that serve people who are immunocompromised or at high risk for severe disease to ensure they know how to get rapid testing ● Ensure access and equity in vaccination, testing, treatment, community outreach, support services for disproportionately affected population |
| <p style="text-align: center;">Medium</p> | <ul style="list-style-type: none"> ● If you are immunocompromised or high risk for severe disease <ul style="list-style-type: none"> ○ Talk to your healthcare provider about whether you need to wear a mask and take other precautions (e.g., testing) ○ Have a plan for rapid testing if needed (e.g., having home tests or access to testing) ○ Talk to your healthcare provider about whether you are a candidate for treatments like oral antivirals, PrEP, and monoclonal antibodies ● If you have household or social contact with someone at high risk for severe disease <ul style="list-style-type: none"> ○ consider self-testing to detect infection before contact ○ consider wearing a mask when indoors with them ● Stay up to date with COVID-19 vaccines and boosters ● Maintain improved ventilation throughout indoor spaces when possible ● Follow CDC recommendations for isolation and quarantine, including getting tested if you are exposed to COVID-19 or have symptoms of COVID-19 | <ul style="list-style-type: none"> ● Protect people at high risk for severe illness or death by ensuring equitable access to vaccination, testing, treatment, support services, and information ● Consider implementing screening testing or other testing strategies for people who are exposed to COVID-19 in workplaces, schools, or other community settings as appropriate ● Implement enhanced prevention measures in high-risk congregate settings (see guidance for correctional facilities and homeless shelters) ● Distribute and administer vaccines to achieve high community vaccination coverage and ensure health equity ● Maintain improved ventilation in public indoor spaces ● Ensure access to testing, including through point-of-care and at-home tests for all people <ul style="list-style-type: none"> ○ Communicate with organizations and places that serve people who are immunocompromised or at high risk for severe disease to ensure they know how to get rapid testing ● Ensure access and equity in vaccination, testing, treatment, community outreach, support services for disproportionately affected populations |

High

- Wear a well-fitting mask indoors in public, regardless of vaccination status (including in K-12 schools and other indoor community settings)
- If you are immunocompromised or high risk for severe disease
 - Wear a mask or respirator that provides you with greater protection
 - Consider avoiding non-essential indoor activities in public where you could be exposed
 - Talk to your healthcare provider about whether you need to take other precautions (e.g., testing)
 - Have a plan for rapid testing if needed (e.g., having home tests or access to testing)
 - Talk to your healthcare provider about whether you are a candidate for treatments like oral antivirals, PrEP, and monoclonal antibodies
- If you have household or social contact with someone at high risk for severe disease
 - consider self-testing to detect infection before contact
 - consider wearing a mask when indoors with them
- Stay up to date with COVID-19 vaccines and boosters
- Maintain improved ventilation throughout indoor spaces when possible
- Follow CDC recommendations for isolation and quarantine, including getting tested if you are exposed to COVID-19 or have symptoms of COVID-19
- Consider setting-specific recommendations for prevention strategies based on local factors
- Implement healthcare surge support as needed
- Protect people at high risk for severe illness or death by ensuring equitable access to vaccination, testing, treatment, support services, and information
- Consider implementing screening testing or other testing strategies for people who are exposed to COVID-19 in workplaces, schools, or other community settings as appropriate
- Implement enhanced prevention measures in high-risk congregate settings (see guidance for correctional facilities and homeless shelters)
- Distribute and administer vaccines to achieve high community vaccination coverage and ensure health equity
- Maintain improved ventilation in public indoor spaces
- Ensure access to testing, including through point-of-care and at-home tests for all people
 - Communicate with organizations and places that serve people who are immunocompromised or at high risk for severe disease to ensure they know how to get rapid testing
- Ensure access and equity in vaccination, testing, treatment, community outreach, support services for disproportionately affected populations